

# Sustained efficacy and safety of bimekizumab in patients with active psoriatic arthritis and prior inadequate response to tumor necrosis factor inhibitors: Results from the Phase 3 BE COMPLETE study and its open-label extension up to 1 year

## Objective

To assess the long-term efficacy and safety of bimekizumab (BKZ) treatment up to 52 weeks in patients with active psoriatic arthritis (PsA) and prior inadequate response or intolerance to tumor necrosis factor- $\alpha$  inhibitors (TNFi-IR).

## Introduction

- BKZ is a humanized monoclonal IgG1 antibody that selectively inhibits interleukin (IL)-17 in addition to IL-17A.
- BKZ has shown superior efficacy to 16 weeks versus placebo (PBO) and tolerability in patients with active PsA in two phase 3 studies, BE OPTIMAL (naïve to biologic disease-modifying antirheumatic drugs [bDMARDs]) and BE COMPLETE (prior inadequate response or intolerance to TNFi-IR).<sup>1,2</sup>
- The efficacy and tolerability of BKZ to 52 weeks has also been demonstrated in BE OPTIMAL.<sup>3</sup>
- Patients with PsA and TNFi-IR typically exhibit reduced treatment responses compared with biologic-naïve patients,<sup>4,5</sup> so identifying treatments that effectively manage the long-term clinical needs of these patients is important.

## Methods

- BE COMPLETE (NCT03896581) included a 16-week double-blind, PBO-controlled period.<sup>2</sup>
- Patients were randomized 2:1 to subcutaneous BKZ 160 mg or PBO every 4 weeks (Q4W).
- Patients who completed Week 16 were eligible for entry into an open-label extension, BE VITAL (NCT04009499; Figure 1). Upon entry, PBO-randomized patients switched to receive BKZ (PBO/BKZ).
- BE VITAL included patients from BE OPTIMAL and BE COMPLETE; data here are only for patients randomized at baseline (Week 0) of BE COMPLETE, up to 1 year.
- Efficacy data reported are observed case or have imputed missing data using non-responder imputation (binary) or multiple imputation (continuous).
- The number of treatment-emergent adverse events (TEAEs) to Week 52 are reported for patients who received  $\geq 1$  dose of BKZ, including patients who switched from PBO to BKZ at Week 16.


## Results

- 388/400 (97.0%) patients completed Week 16; 377 (94.3%) entered BE VITAL and 347 (86.8%) completed Week 52.
- Baseline characteristics were comparable between groups (Table 1).
- Improvements in joint and skin responses with BKZ treatment at Week 16 were sustained to Week 52 (Figure 2 and Table 2).
- Patients who switched to BKZ at Week 16 demonstrated improvements in efficacy responses to Week 52 (Figure 2 and Table 2).
- To Week 52, 243/388 (62.6%) patients had  $\geq 1$  TEAE whilst receiving BKZ (exposure-adjusted incident rate [EAIR]: 126.0 per 100 patient-years; Table 3).
- The most frequent TEAEs were coronavirus infection, oral candidiasis, nasopharyngitis and urinary tract infection (Table 3).
- All *Candida* infections were mild or moderate and none were systemic.
- Two cases of oral candidiasis led to study discontinuation.
- There was one death, considered unrelated to study treatment by the investigator (BKZ-treated patient with a history of cardiac events).

## Conclusions

In patients with PsA and prior TNFi-IR, bimekizumab treatment demonstrated sustained improvements across joints and skin from Week 16 to Week 52. Patients who switched to bimekizumab at Week 16 also displayed meaningful improvements in efficacy responses at Week 52. The safety profile was consistent with previous reports.<sup>1-3</sup>

Figure 1 BE COMPLETE and BE VITAL study design



\*BE VITAL includes patients from the BE OPTIMAL and BE COMPLETE studies; results are only presented for patients from BE COMPLETE. BKZ-treated patients were eligible to receive rescue therapy from Week 16 at the discretion of the investigator, while continuing to receive BKZ.

Table 1 Baseline patient demographics and disease characteristics

	PBO n=133	BKZ 160 mg Q4W n=267
Age, years, mean (SD)	51.3 (12.9)	50.1 (12.4)
Male, n (%)	60 (45.1)	130 (48.7)
BMI, kg/m <sup>2</sup> , mean (SD)	29.0 (5.4)	30.1 (6.5)
Time since first diagnosis of PsA, <sup>a</sup> years, mean (SD)	9.2 (8.1)	9.6 (9.9)
TJC (of 68 joints), mean (SD)	19.3 (14.2)	18.4 (13.5)
SJC (of 66 joints), mean (SD)	10.3 (8.2)	9.7 (7.5)
hs-CRP $\geq 6$ mg/L, n (%)	59 (44.4)	118 (44.2)
Patients with psoriasis involving $\geq 3\%$ BSA, n (%) / PASI score, <sup>b</sup> mean (SD)	88 (66.2) / 8.5 (6.6)	176 (65.9) / 10.1 (9.1)
HAQ-DI score, mean (SD)	1.04 (0.69)	0.97 (0.59)
Dactylitis (LDI >0), <sup>c,d</sup> n (%) / LDI score, <sup>c</sup> mean (SD)	14 (10.5) / 66.4 (127.6)	34 (12.7) / 72.7 (114.4)
Enthesitis (LEI >0), <sup>c,e</sup> n (%) / LEI score, <sup>e</sup> mean (SD)	36 (27.1) / 2.9 (1.6)	106 (39.7) / 2.6 (1.5)
Nail psoriasis (mNAPSI >0), <sup>d</sup> n (%) / mNAPSI score, <sup>e</sup> mean (SD)	83 (62.4) / 4.5 (2.8)	159 (59.6) / 4.3 (2.8)


Randomized set. \*Data missing for 1 PBO patient; 1 BKZ patient; <sup>a</sup>Patients with psoriasis involving  $\geq 3\%$  BSA at baseline; <sup>b</sup>The presence of dactylitis was defined by a score greater than 0 on the Leeds Dactylitis Index (higher scores indicate a greater number of affected sites); <sup>c</sup>Data missing for 1 PBO patient; <sup>d</sup>In patients with dactylitis at baseline; <sup>e</sup>The presence of enthesitis was defined by a score greater than 0 on the Leeds Enthesitis Index (range 0 to 6, with higher scores indicating a greater number of affected sites). <sup>f</sup>n patients with enthesitis at baseline; <sup>g</sup>n patients with nail psoriasis at baseline.

ACR: American College of Rheumatology; ACR20/50/70:  $\geq 20/50/70\%$  improvement from baseline in ACR criteria; AE: adverse event; bDMARD: biologic disease-modifying antirheumatic drug; BKZ: bimekizumab; BMI: body mass index; BSA: body surface area; CFb: change from baseline; EAIR: exposure-adjusted incident rate per 100 patient-years; HAQ-DI: Health Assessment Questionnaire-Disability Index; hs-CRP: high-sensitivity C-reactive protein; IBD: inflammatory bowel disease; IL: interleukin; LDI: Leeds Dactylitis Index; LEI: Leeds Enthesitis Index; MACE: major adverse cardiovascular event; MDA: minimal disease activity; MI: multiple imputation; mNAPSI: modified Nail Psoriasis Severity Index; NRI: non-responder imputation; OC: observed case; OLE: open-label extension; PASI: Psoriasis Area and Severity Index; PBO: placebo; PsA: psoriatic arthritis; PsAID-12: Psoriatic Arthritis Impact of Disease-12; PYAR: patient years at risk; Q4W: every four weeks; SD: standard deviation; SE: standard error; SF-36 PCS: Short-Form 36-item Health Survey Physical Component Summary; SJC: swollen joint count; TEAE: treatment-emergent adverse event; TJC: tender joint count; TNFi: tumour necrosis factor- $\alpha$  inhibitor; TNFi-IR: prior inadequate response or intolerance to tumor necrosis factor- $\alpha$  inhibitors.

Affiliations: <sup>1</sup>Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Diseases, University of Oxford and Oxford Biomedical Research Centre, Oxford University Hospitals NHS Trust, Oxford, UK; <sup>2</sup>Amsterdam Rheumatology & Clinical Immunology Center, Amsterdam, and Zuyderland MC, Heerlen, The Netherlands; <sup>3</sup>College of Medical Veterinary and Life Sciences, University of Glasgow, Glasgow, UK; <sup>4</sup>Swedish Medical Center and Providence St. Joseph Health and University of Washington, Seattle, Washington, USA; <sup>5</sup>Allergy, Immunology & Rheumatology Division, University of Rochester Medical School, Rochester, New York, USA; <sup>6</sup>The First Department of Internal Medicine, University of Occupational and Environmental Health, Japan, Kitakyushu, Fukuoka, Japan; <sup>7</sup>Department of Dermatology, The Jikei University School of Medicine, Tokyo, Japan; <sup>8</sup>Division of Rheumatology, University Hospital and Fraunhofer Institute for Translational Medicine & Pharmacology ITMP, Fraunhofer Center of Excellence Immune-Mediated Diseases CIMD, Frankfurt, Goethe University, Frankfurt am Main, Germany; <sup>9</sup>Schroeder Arthritis Institute, Krembil Research Institute, Toronto Western Hospital, University Health Network, University of Toronto, Ontario, Canada; <sup>10</sup>Sorbonne Université and Pitié-Salpêtrière Hospital, Paris, France; <sup>11</sup>Department of Dermatology, The Icahn School of Medicine at Mount Sinai, New York, New York, USA; <sup>12</sup>Dermatology Center, Northern Care Alliance NHS Foundation Trust; <sup>13</sup>NHR Manchester Biomedical Research Centre, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK; <sup>14</sup>UCB Pharma, Slough, UK; <sup>15</sup>UCB Pharma, Morrisville, North Carolina, USA; <sup>16</sup>Department of Dermatology, Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts, USA; <sup>17</sup>Division of Rheumatology, Department of Medicine, Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts, USA.

References: McInnes IB, et al. Lancet 2023;401:25-37; Merola JF, et al. Ann Rheum Dis 2022;71:2840-47; Ritchlin C, et al. JAMA Dermatol 2022;158:1627-35. Author Contributions: Substantial contributions to study conception/design, or acquisition/analysis/interpretation of data: LCC, RBML, IBM, PJM, CTR, YF, AA, FB, DDG, LG, ABG, RBW, BI, RB, JC, JFM. Author Disclosures: LCC: Grants/research support from AbbVie, Amgen, Celgene, Eli Lilly, Gilead, Janssen, Novartis, Pfizer, and UCB Pharma; Consultant for AbbVie, Amgen, Celgene, Eisai, Eli Lilly, Gilead, Janssen, Novartis, Pfizer, and UCB Pharma; Consultant for AbbVie, Amgen, Celgene, Eisai, Eli Lilly, Gilead, Janssen, Novartis, Pfizer, and UCB Pharma; Owner of Rheumatology Consultancy BV, an AMS company under Dutch law; IBM: Consulting fees and honoraria from AbbVie, AstraZeneca, BMS, Eli Lilly, Novartis, Pfizer, and UCB Pharma; Takeda, UCB Pharma, and Ventyx; Speakers' bureau for AbbVie, Amgen, Eli Lilly, Janssen, Novartis, Pfizer, and UCB Pharma; Research support from BMS, Boehringer Ingelheim, Celgene, Janssen, Novartis, and UCB Pharma; PJM: Research grants from AbbVie, Acelyrin, Amgen, BMS, Eli Lilly, Gilead, Janssen, Novartis, Pfizer, Sun Pharma, and UCB Pharma; Consultant fees from AbbVie, Acelyrin, Amgen, BMS, Boehringer Ingelheim, Eli Lilly, Gilead, Janssen, Novartis, Pfizer, and UCB Pharma; Received grants from Chugai, Eisai, Mitsubishi-Tanabe, and Taisho; AA: Honoraria and/or research grants from AbbVie, Amgen, BMS, Boehringer Ingelheim, Eisai, Eli Lilly, Janssen, Kyowa Kirin, LEO Pharma, Maruho, Mitsubishi Tanabe Pharma, Novartis, Pfizer, Sun Pharma, Taiso Pharma, Torii Pharmaceutical Co., and UCB Pharma; FB: Consultant and/or speaker and/or investigator for AbbVie, Amgen, Boehringer Ingelheim, Celgene, Chugai, Eli Lilly, Genzyme, GSK, Janssen, MSD, Moonlake Pharma, Novartis, Pfizer, Roche, Sandoz, and Sanofi; DDG: Consultant and/or research grant support from AbbVie, Amgen, BMS, Celgene, Eli Lilly, Galapagos, Gilead, Janssen, Novartis, Pfizer, and UCB Pharma; LG: Grants from AbbVie, Biogen, Eli Lilly, Novartis, Sandoz, and UCB Pharma; Personal fees from AbbVie, Amgen, BMS, Celgene, Eli Lilly, Galapagos, Janssen, MSD, Novartis, Pfizer, Sandoz, and UCB Pharma; ABG: Received honoraria as an advisory board member and consultant for Amgen, AnaptysBio, Avotres Therapeutics, BMS, Boehringer Ingelheim, Dice Therapeutics, Eli Lilly, Janssen, Novartis, Sanofi, UCB Pharma, and Xbiotech; Received research/educational grants from AnaptysBio, BMS, Moonlake Immunotherapeutics, Novartis, and UCB Pharma (all paid to Mount Sinai School of Medicine); RBW: Consulting fees from AbbVie, Amgen, BMS, Dermavant, Eli Lilly, Janssen, LEO Pharma, Pfizer, Novartis, Regeneron, Sanofi, Sun Pharma, and UCB Pharma; Acknowledgments: We would like to thank the patients and their caregivers in addition to all the investigators and their teams who contributed to this study. The authors acknowledge Heather Edens, PhD, UCB Pharma, Smyrna, Georgia, USA, for publication coordination; Nadine Goldammer, PhD, UCB Pharma, Monheim, Germany, for their work as clinical program delivery lead for the bimekizumab PsA program; David Morgan, PhD, Costello Medical, Manchester, UK for medical writing and editorial assistance, and the Costello Medical Creative team for design support. This study was funded by UCB Pharma. All costs associated with development of this presentation were funded by UCB Pharma.

Figure 2 ACR, PASI and MDA response rates over time to Week 52 (NRI and OC)



Randomized set. For binary variables, p values were calculated using a logistic regression model with treatment, prior TNFi exposure, and region as stratification factors. Nominal p values are not powered or adjusted for multiplicity and should not be used to assess statistical significance. N values are NRI/OC at Week 0 and OC at Week 16 and Week 52. \*In patients with psoriasis involving  $\geq 3\%$  BSA at baseline.

Table 2 Additional efficacy endpoints at Week 16 and Week 52 (NRI)

	PBO (Weeks 0-16) n=133		BKZ 160 mg Q4W (Weeks 16-52) n=133	
	Week 16	Week 52	Week 16	Week 52
PASI75 response <sup>a</sup>	9/88 (10.2)	71/88 (80.7)	145/176 (82.4)	148/176 (84.1)
Enthesitis resolution <sup>b</sup>	8/36 (22.2)	21/36 (58.3)	52/106 (49.1)	60/106 (56.6)
Dactylitis resolution <sup>c</sup>	6/14 (42.9)	12/14 (85.7)	24/34 (70.6)	29/34 (85.3)
Nail psoriasis resolution <sup>d</sup>	12/83 (14.5)	51/83 (61.4)	73/159 (45.9)	107/159 (67.3)
HAQ-DI CFb, MI, mean (SE)	-0.07 (0.04)	-0.35 (0.05)	-0.38 (0.03)	-0.39 (0.03)
	Week 16	Week 40*	Week 16	Week 40*
SF-36 PCS CFb, MI, mean (SE)	1.4 (0.7)	7.3 (0.9)	7.3 (0.5)	8.4 (0.6)

Randomized set. Previously reported data through Week 16 included for reference. \*Data not collected at Week 52 for SF-36 PCS. <sup>a</sup>In patients with psoriasis involving  $\geq 3\%$  BSA at baseline; <sup>b</sup>Patients with enthesitis at baseline (LEI >0); <sup>c</sup>Patients with dactylitis at baseline (LDI >0); <sup>d</sup>Patients with nail psoriasis at baseline (mNAPSI >0).

Laura C. Coates,<sup>1</sup> Robert B.M. Landewé,<sup>2</sup> Iain B. McInnes,<sup>3</sup> Philip J. Mease,<sup>4</sup> Christopher T. Ritchlin,<sup>5</sup> Yoshiya Tanaka,<sup>6</sup> Akihiko Asahina,<sup>7</sup> Frank Behrens,<sup>8</sup> Dafna D. Gladman,<sup>9</sup> Laure Gossec,<sup>10</sup> Alice B. Gottlieb,<sup>11</sup> Richard B. Warren,<sup>12,13</sup> Barbara Ink,<sup>14</sup> Rajan Bajracharya,<sup>14</sup> Jason Coarse,<sup>15</sup> Joseph F. Merola<sup>16,17</sup>