

Patient and Caregiver Perceptions of Acute Seizure Medications and the Rapid and Early Seizure Termination (REST) Approach: Qualitative Interviews

1. UCB, Slough, UK
2. Kielo Research, Zug, Switzerland
3. Centennial Children's Hospital, Nashville, TN, USA
4. Vall d'Hebron University Hospital, Barcelona, Spain
5. Montefiore Medical Center, Bronx, NY, USA
6. Patient Author, Chicago, IL, USA
7. Kielo Research, York, UK
8. UCB, Brussels, Belgium

Kerrie-Anne Ho¹
 Tommi Tervonen²
 Jesus Eric Pina-Garza³
 Manuel Toledo Argany⁴
 Sheryl R Haut⁵
 Tanya Bhatia⁶
 Cecilia Jimenez-Moreno⁷
 Cédric Lalouaux⁸

P4-003

Background

- Rapid and Early Seizure Termination (REST) is essential to prevent seizures becoming prolonged seizures (PS) or progressing in severity.¹
- REST is a new management paradigm that encompasses the acute treatment of ongoing seizures, with the aim of being rapid (quick onset of action) and early (used as close to seizure onset as possible).¹
- However, among people with epilepsy (PwE) and caregivers, awareness of the REST paradigm and perceptions around acute (on-demand) medications are not well known.

Objective

- To understand the experience of PS and acute medications among PwE and caregivers of PwE and assess the understanding and perceptions of the REST paradigm via qualitative interviews.

Methods

- Participants included PwE aged ≥18 years, with a diagnosis of epilepsy or an epilepsy syndrome, and who had experienced ≥1 PS in the prior 12 months (defined as either a tonic-clonic or motor type seizure [lasting ≥2 minutes]; a non-motor seizure with impaired awareness or an absence seizure [lasting ≥2 minutes]; or a non-motor seizure when still aware [lasting ≥5 minutes]), and caregivers (aged ≥18 years) of PwE (aged ≥12 years) from France, Italy, Poland, Spain, the United Kingdom (UK), and the United States (US).
- Participants took part in 60- to 90-minute qualitative interviews regarding experiences with auras, epilepsy concepts, experience and unmet needs of acute medications, and perceptions of the REST paradigm.
- PwE answered questions/gave feedback regarding their own experiences; caregivers' experiences/observations were their own and caregivers were not answering on behalf of PwE.
- Interviews were recorded and analyzed using formal qualitative coding techniques.

Results

PARTICIPANTS' CHARACTERISTICS AND SEIZURE EXPERIENCE

- 53 participants took part in qualitative interviews: 18 PwE aged ≥18 years, 18 caregivers of adult PwE (aged ≥18 years), and 17 caregivers of adolescent PwE (aged ≥12 to <17 years).
- 83% of participants were female; median (25th percentile [Q1], 75th percentile [Q3]) age was 47 (40, 51) years.
- Nine participants each were from France, Italy, Poland, the UK, and the US; 8 were from Spain.

Sociodemographic characteristics and seizure history

| | Overall (N=53) | Adult PwE (aged ≥18 years) (N=18) | Caregiver of adult PwE (aged ≥18 years) (N=18) | Caregiver of adolescent PwE (aged ≥12 to <17 years) (N=17) |
|--|----------------|-----------------------------------|--|--|
| Age of participant, median (Q1, Q3), years | 47 (40, 51) | 47 (30, 54) | 50 (42, 56) | 46 (39, 50) |
| Female, n (%) | 44 (83) | 11 (61) | 17 (94) | 16 (94) |
| Most common seizure types,^a n (%) | | | | |
| Generalized or bilateral seizures/tonic-clonic | 35 (66) | 9 (50) | 13 (72) | 13 (76) |
| Focal seizures without loss of awareness | 22 (42) | 7 (39) | 8 (44) | 7 (41) |
| Focal seizures with loss of awareness | 20 (38) | 7 (39) | 8 (44) | 5 (29) |
| Generalized or bilateral seizures non-motor | 13 (25) | 6 (33) | 3 (17) | 4 (24) |
| Unknown onset with present visible physical movement, jerks, or body stiffness | 6 (11) | 2 (11) | 3 (17) | 1 (6) |
| Unknown onset without any visible movements of the body or as an absence | 3 (6) | 1 (6) | 1 (6) | 1 (6) |
| Only experiences tonic-clonic seizures | 14 (26) | 2 (11) | 5 (28) | 7 (41) |
| Multiple seizure types | 27 (51) | 9 (50) | 11 (61) | 7 (41) |
| Average number of seizures per month during the prior 12 months, n (%) | | | | |
| <3 | 12 (23) | 6 (33) | 2 (11) | 4 (24) |
| 3-5 | 17 (32) | 6 (33) | 6 (33) | 5 (29) |
| 6-10 | 6 (11) | 2 (11) | 3 (17) | 1 (6) |
| 11-20 | 5 (9) | 2 (11) | 2 (11) | 1 (6) |
| >20 | 13 (25) | 2 (11) | 5 (28) | 6 (35) |
| Average duration of most common seizure experienced during the prior 12 months, n (%) | | | | |
| <1 minute | 4 (8) | 2 (11) | 1 (6) | 1 (6) |
| 1-2 minutes | 16 (30) | 5 (28) | 7 (39) | 4 (24) |
| 2-4 minutes | 21 (40) | 6 (33) | 6 (33) | 9 (53) |
| 4-5 minutes | 6 (11) | 3 (17) | 2 (11) | 1 (6) |
| >5 minutes | 6 (11) | 2 (11) | 2 (11) | 2 (12) |

- For caregivers (n=35), median (Q1, Q3) age of the PwE being cared for was 19 (15, 28) years.
- 71% of caregivers were parents and 6% had been caring for a PwE for >10 years.

Overview

QUESTION


Among people with epilepsy (PwE) and caregivers of PwE, what is the experience of prolonged seizures (PS) and acute medications and the level of understanding and perception of the Rapid and Early Seizure Termination (REST) paradigm?

INVESTIGATION


Participants included PwE aged ≥18 years, with a diagnosis of epilepsy/epilepsy syndrome who had experienced ≥1 PS (≥2 minutes) in the prior 12 months and caregivers (aged ≥18 years) of PwE (aged ≥12 years). Participants took part in qualitative interviews regarding experiences with auras, epilepsy concepts, experience and unmet needs of acute medications, and perceptions of the REST paradigm.

RESULTS


Experience with auras




Knowledge of epilepsy concepts



Current treatments and experience with acute medications



Perceptions of the REST paradigm




CONCLUSIONS


Many participants reported they could confidently predict seizures based on auras. Most participants reported using or administering a benzodiazepine as an acute medication for PS. Most participants felt that REST is a feasible approach and suggested it could provide relief with being able to act earlier before seizures progress. REST could be a beneficial and fast-acting approach to prevent seizures becoming PS or progressing in severity.

EXPERIENCE WITH AURAS


Aura or early symptoms at the very beginning of seizures^a



Confidence regarding aura or warning signs progressing to a seizure




Confidence recognizing an upcoming seizure as prolonged from experience with auras



KNOWLEDGE OF EPILEPSY CONCEPTS

- A specific time frame for the definition of a PS was given by 74% of participants.


Definitions of PS




CURRENT TREATMENTS AND UNMET NEEDS

- 87% of participants used or administered a benzodiazepine as an acute medication for PS.


Modes of administration of current acute medications used to treat PS^a




Main expectations/goals when using acute medication (N=53)^a



Experience with acute medication




Changes participants would like to see regarding acute medication (N=53)



PERCEPTIONS OF THE REST PARADIGM

Expectations regarding time for a REST medication to act (N=53)




- 53% of participants (28/53) perceived a REST medication as quick/fast if it acts in ≤1 minute.


Perceptions of the REST paradigm

- The concept of REST (ie, treating seizures early with a fast-acting treatment) was explained to participants, and 15% (n=8/53) indicated they were already treating seizures early.
- 45 participants were not treating seizures early, of whom 67% (n=30/45) perceived REST to be a feasible approach.

Most mentioned perceived advantages regarding REST (N=53)^a



Most mentioned perceived concerns regarding REST (N=53)^a



Conclusions

- Many participants reported they could confidently predict seizures based on auras.
- Most participants reported using or administering a benzodiazepine as acute medication for PS.
- Unmet needs of current acute medication were effectiveness (being more effective and/or quicker to act) and mode of administration.
- Most participants felt that REST is a feasible approach and suggested it could provide relief with being able to act earlier before seizures progress.
- This study highlights the unmet need for REST therapies; REST could be a beneficial and fast-acting approach to prevent seizures becoming PS or progressing in severity.

References

- Pina-Garza JE. *Epileptic Disord* 2024;26(4):484-497.
- Beniczky S, et al. *Epilepsia* 2025;66(6):1804-1823.

UCB-sponsored. UCB was involved in the design of the study, the collection, analysis, and interpretation of data, and review of the poster. The authors thank the patients and their caregivers in addition to the investigators and their teams who contributed to this study. The authors acknowledge Tom Grant, PhD (UCB, Slough, UK) for managing the development of the poster, and Emma Buttl, PhD (Envision Spark, an Envision Medical Communications agency, a part of Envision Pharma Group, Horsham, UK) for writing assistance, which was funded by UCB. Author contributions: All authors contributed to designing the study, analyzing and interpreting the data, critically reviewing the poster, and approving the final version for presentation. Author disclosures: K.A.Ho and C.Lalouaux are employees of UCB and receive stock or stock options from their employer. T.Tervonen and C.Jimenez-Moreno are employees of Kielo Research, who received funds from UCB related to this study. T.Tervonen holds stock in Kielo Research. J.E.Pina-Garza has received royalties from Elsevier; received consulting fees as an advisor to Eisai, Jazz, Marinus, Neurelis, SK Life Science, UCB, and Zogenix; and received payment or honoraria for speaker bureaus for Eisai, Jazz, Marinus, Neurelis, SK Life Science, UCB, and Zogenix. M.Toledo Argany declares consulting honoraria and research fundings from Angelini, Arvelle, Bial, Eisai, Esteve, GW Pharmaceuticals, Jazz, Neuropharm, Neuroelectrics, Takeda, and UCB. SR.Haut is a consultant for Neurelis, UCB, and Ventus; and a data safety monitoring board member for AbbVie. T.Bhatia did not report any disclosures.

For a copy of this poster, use your smartphone to scan the QR code or contact UCB-CARES[®]
 Phone: +1-844-599-CARE (2273)
 Email: UCB-CARES@ucb.com

American Academy of Neurology 78th Annual Meeting
 Chicago, IL, USA | April 18-22, 2026
 Previously presented at American Epilepsy Society 79th Annual Meeting, Atlanta, GA, USA, December 5-9, 2025