

Bimekizumab Efficacy by Patient Subgroups in Moderate to Severe Hidradenitis Suppurativa: 3-Year Phase 3 Results from BE HEARD EXT

Christopher J. Sayed,^{1,2} Martina L. Porter,³ Alejandro Molina-Leyva,^{2,4} Jacek C. Szepietowski,^{2,5} Yosuke Ishitsuka,⁶ Evelina van Ree,⁷ Christina Crater,⁸ Tom Vaux,⁸ Georgios Kokolakis^{2,9}

¹Department of Dermatology, University of North Carolina School of Medicine, Chapel Hill, NC, USA; ²European Hidradenitis Suppurativa Foundation (EHSF) e.V., Dessau, Germany; ³Beth Israel Deaconess Medical Center, Department of Dermatology, Harvard Medical School, Boston, MA, USA; ⁴Dermatology Department, Hospital Universitario Virgen de las Nieves - Ibs. Granada, Granada, Spain; ⁵Division of Dermatology, Venereology and Clinical Immunology, Faculty of Medicine, Wrocław University of Science and Technology, Wrocław, Poland; ⁶Department of Dermatology, Osaka University Graduate School of Medicine, Osaka, Japan; ⁷UCB, Breda, The Netherlands; ⁸UCB, Morrisville, NC, USA; ⁹Department of Dermatology, Venereology and Allergology, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany.

Objective

To report bimekizumab (BKZ) efficacy and maintenance of clinical response over 3 years across different subgroups of patients with moderate to severe hidradenitis suppurativa (HS) using phase 3 (BE HEARD I&II) and open label extension (BE HEARD EXT) data.

Background

- HS disease varies between individuals based on patient and disease characteristics, such as sex and disease duration.¹
- Biological therapies which provide consistent efficacy, regardless of patient characteristics, are needed.
- BKZ is a humanized IgG1 monoclonal antibody that selectively inhibits interleukin (IL)-17A and IL-17F.²

Methods

- Data were pooled from the phase 3 BE HEARD I&II trials and their open-label extension, BE HEARD EXT.^{3,4}
- Patients completing the 48-week BE HEARD I&II studies could enroll in BE HEARD EXT and receive open-label BKZ 320 mg every 2 weeks (Q2W) or every 4 weeks (Q4W) based on HS Clinical Response (HiSCR)-90 response, averaged from Weeks 36, 40, and 44.
- We report the rates of patients achieving HiSCR50/90 for subgroups at Years 1, 2, and 3 (Weeks 48, 96, and 148).
- Data are reported as observed case (OC) for patients randomized to BKZ 320 mg from baseline in BE HEARD I&II who entered BE HEARD EXT and continued to receive BKZ (BKZ Total).

Results

- 657 BE HEARD I&II completers entered BE HEARD EXT; 556 received BKZ from baseline.
- Baseline characteristics and clinical characteristics of patients are reported in the **Table**.
- Most patients across subgroups achieved HiSCR50, with responses maintained through to Year 3 (**Figure 1**).
- Similarly, most patients achieved HiSCR90 at Year 2, with responses maintained through to Year 3, indicating that deeper responses take longer to achieve (**Figure 2**).

Conclusions

Patients treated with bimekizumab demonstrated high clinical response rates across all subgroups at Year 1, with responses maintained through to Year 3. These results emphasize the long-term benefit of bimekizumab as an effective treatment option for patients with hidradenitis suppurativa, regardless of patient demographics and disease characteristics.

Summary




Table Baseline demographics

	BKZ Total ^a N=556
Age, years, mean (SD)	36.3 (12.2)
Sex, female, n (%)	299 (53.8)
Racial group, White, n (%)	448 (80.6)
Weight, kg, mean (SD)	96.2 (23.5)
BMI, kg/m ² , mean (SD)	32.5 (7.8)
Duration of HS, years, mean (SD)	7.4 (7.1)
Hurley stage, n (%)	
II	303 (54.5)
III	253 (45.5)
Prior biologic use ^b , n (%)	112 (20.1)
IHS4 severity, n (%)	
Mild: ≤3	0
Moderate: 4–10	70 (12.6)
Severe: ≥11	486 (87.4)

OLE set: N=657; included only patients who entered BE HEARD EXT at Week 48. [a] BKZ Total comprised patients randomized to BKZ from baseline in BE HEARD I&II who entered BE HEARD EXT and continued to receive BKZ; [b] Patients received prior biologic therapy for any indication.

BKZ: bimekizumab; **BMI:** body mass index; **HiSCR50/90:** ≥50/90% reduction from baseline in the total abscess and inflammatory nodule (AN) count, with no increase from baseline in abscess or draining tunnel (DT) count; **HS:** hidradenitis suppurativa; **IHS4:** International Hidradenitis Suppurativa Severity Score System; **IL:** Interleukin; **OC:** observed case; **OLE:** open-label extension; **Q2W:** every 2 weeks; **Q4W:** every 4 weeks; **SD:** standard deviation.

References: ¹Schrader AMR, et al. J Am Acad Dermatol 2014;71:460–7. ²Adams R, et al. Front Immunol 2020;11:1894; ³Kimball AB, et al. Lancet 2024;403:2504–19 (NCT04242446, NCT04242498); ⁴BE HEARD EXT (NCT04901195); www.clinicaltrials.gov/study/NCT04901195. **Author Contributions:** Substantial contributions to study conception/design, or acquisition/analysis/interpretation of data: **CJS, MLP, AM-L, JCS, YI, EVR, CC, TV, GK**; Drafting of the publication, or reviewing it critically for important intellectual content: **CJS, MLP, AM-L, JCS, YI, EVR, CC, TV, GK**; Final approval of the publication: **CJS, MLP, AM-L, JCS, YI, EVR, CC, TV, GK**. **Author Disclosures:** **CJS:** Investigator for AbbVie, AstraZeneca, ChemoCentryx, Incyte, InflaRx, Novartis, and UCB; consultancy fees from AbbVie, Alumis, Arcutis, AstraZeneca, InflaRx, Incyte, Logical Images, MoonLake Immunotherapeutics, Oruka, Sandoz, Sanofi, Sonoma Biotherapeutics, UCB, Ventyx, and Zura Bio; consultant for Almirall, FIDE, Insmed, Oruka, Otsuka, Propeller Biosciences, Spyre, and Trifecta Clinical/WCG; investigator for AnaptysBio, Bayer, Bristol Myers Squibb, OASIS Pharmaceuticals, and Regeneron; received royalties from Beth Israel Deaconess Medical Center; received fellowship funding to institution from AbbVie. **AM-L:** Received support for attending meetings, served as an advisory board member, and participated in clinical trials from AbbVie, Almirall, Incyte, MoonLake Immunotherapeutics, MSD, Novartis, Sanofi, and UCB. **JCS:** Consultant and advisory board member of AbbVie, Almirall, Boehringer Ingelheim, Galderma, LEO Pharma, Novartis, Pierre Fabre, Sandoz, Sanofi Genzyme, and UCB; speaker for AbbVie, Almirall, Boehringer Ingelheim, Eli Lilly and Company, Janssen, LEO Pharma, Novartis, Pfizer, Pierre Fabre, Sanofi Genzyme, and UCB; investigator for AbbVie, Acelyrin, Almirall Hermal, Amgen, AnaptysBio, Argenx, Aslan, Boehringer Ingelheim, Biocom, Bio Thera, Bristol Myers Squibb, Celltrion, CuraTeQ Biologics, DICE Therapeutics, Eli Lilly and Company, Galapagos, Galderma, Helm AG, Incyte, InflaRx, Janssen, Kiniksa, Kymab Limited, LEO Pharma, Medimmune, Menlo Therapeutics, Merck, MoonLake Immunotherapeutics, Novartis, Pierre Fabre, Pfizer, Regeneron Pharmaceuticals, Takeda, Teva, Trevi Therapeutics, UCB, Uni Therapeutics, and Ventyx Bioscience. **YI:** Nothing to disclose. **EVR, CC, TV, GK:** Employees and shareholders of UCB. **AK:** Received travel grants or honoraria, has been a consultant member of advisory boards and speaker bureaus, or has served as an investigator for AbbVie, Actelion, Almirall, Amgen, Basilea, Biogen, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Eli Lilly and Company, Hexal-Sandoz, Janssen, LEO Pharma, L'Oréal, MSD, Novartis, Pfizer, Sanofi, Takeda, and UCB. **Acknowledgments:** We would like to thank the patients and their caregivers in addition to all the investigators and their teams who contributed to these studies. The authors acknowledge Susanne Wiegatz, UCB, Monheim am Rhein, Germany for publication coordination, Ujjawal Kumar, MA, MB, BChir, Costello Medical, Cambridge, UK for medical writing support, Isabel Merrien, PgDip, for editorial assistance, and the Costello Medical Creative team for design support. This study was funded by UCB. All costs associated with development of this poster were funded by UCB.

Figure 1 HiSCR50 achievement across patient subgroups to Year 3





Figure 2 HiSCR90 achievement across patient subgroups to Year 3



To receive a copy of this poster, scan the QR code. Link expiration: August 8, 2026

